



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Beyond the point of no return

Citation for published version:

Buonocore, A, McIntosh, R & Melcher, D 2016, 'Beyond the point of no return: Effects of visual distractors on saccade amplitude and velocity', *Journal of Neurophysiology*, vol. 115, no. 2, pp. 752-762.
<https://doi.org/10.1152/jn.00939.2015>

Digital Object Identifier (DOI):

[10.1152/jn.00939.2015](https://doi.org/10.1152/jn.00939.2015)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of Neurophysiology

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



1 **Beyond the point of no return: effects of visual distractors on**
2 **saccade amplitude and velocity**

3

4

5 Antimo Buonocore¹, Robert D. McIntosh², David Melcher¹

6

7 ¹Center for Mind/Brain Sciences, University of Trento, Italy

8 ²Human Cognitive Neuroscience, University of Edinburgh, United Kingdom

9

10

11

12

13 **Corresponding Author:**

14 Dr Antimo Buonocore

15 Center for Mind/Brain Sciences (CIMEC)

16 University of Trento

17 Corso Bettini, 31

18 38068, Rovereto (TN), Italy

19 Tel: +39 0464 808728

20 antimo.buonocore@unitn.it

21

22 **Abstract**

23 Visual transients, such as a bright flash, reduce the proportion of saccades executed
24 around 60-125 ms after flash onset, a phenomenon known as *saccadic inhibition*.
25 Across three experiments, we apply a similar time-course analysis to the amplitudes
26 and velocities of saccades. Alongside the expected reduction of saccade frequency in
27 the key time period, we report two perturbations of the “main sequence”, one before
28 and one after the period of saccadic inhibition. First, saccades launched between 30 to
29 70 ms following the flash were hypometric, with peak speed exceeding that expected
30 for a saccade of similar amplitude. This finding was in contrast to the common idea
31 that saccades have passed a “point-of-no-return” around 60 ms prior to launching,
32 escaping interference from distractors. The early hypometric saccades observed were
33 not a consequence of spatial averaging between target and distractor locations, as they
34 were found not only following a localized central flash (Experiment 1), but also
35 following a spatially generalized flash (Experiment 2). Second, across experiments,
36 saccades launched at 110 ms post-flash, toward the end of saccadic inhibition, had
37 normal amplitude but a peak speed higher than expected for that amplitude suggesting
38 increased collicular excitation at the time of launching. Overall, the results show that
39 saccades that escape inhibition following a visual transient are not necessarily
40 unaffected, but instead can reveal interference in spatial and kinematic measures.

41

42 **Keywords:** eye movements, saccadic inhibition, main sequence

43 Introduction

44 The main challenge for the oculomotor system in a complex environment is to select
45 when and where to move the eyes in order to land near targets of interest. Saccadic
46 amplitudes range from only a few minutes of arc to over 80° of visual angle, and the
47 kinematics of saccadic performance are generally invariant across tasks and people.
48 Perhaps the best example of this regularity is the lawful monotonic relationship
49 between saccadic amplitude and peak speed, called the *main sequence* (Bahill et al.
50 1975; Collewyn et al. 1988), which holds up to ~60° of visual angle (at which the
51 peak speed saturates at ~500°/s). Another stereotyped feature of the saccadic system is
52 how it responds to sudden transient events. Reingold and Stampe (1999, 2000, 2003,
53 2004) used a highly salient flashed distractor, and revealed a characteristic “dip” in
54 saccadic frequency beginning as early as 60-70 ms after the flash, with maximal
55 depression around 90 ms, rebounding to normal levels by 120-130 ms. This *saccadic*
56 *inhibition* (SI) generalized beyond the text-reading and scene-exploration tasks first
57 tested, with distractors having similar effects in gap, overlap, pro-saccade and anti-
58 saccade tasks (Reingold and Stampe 2002). Interestingly, an analogous effect has
59 been shown for endogenously triggered micro-saccades. “Micro-saccadic inhibition”
60 describes a similar dip in micro-saccade rate, about 100 ms after the presentation of a
61 visual cue (Engbert and Kliegl 2003; Hafed and Clark 2002). As for standard SI, this
62 effect was shown to occur with any sensory transient presented during saccadic
63 planning, supporting the idea of generalized inhibitory mechanisms in the oculomotor
64 system (Hafed and Ignashchenkova 2013).

65 An interesting question concerns the time window during which a saccadic
66 plan is susceptible to interference. The most commonly accepted estimates of *when* a
67 saccade plan can still be modulated have come from double-step tasks, in which

68 participants have to saccade toward a target that sometimes jumps to a second
69 location after initial presentation. The amplitude of the first saccade varies as a
70 function of the delay between the target jump and the onset of the first saccade. When
71 this delay is short, the eye movement will land at the first target location, but for
72 longer delays the saccade will land at the second target. For intermediate delays, the
73 saccade tends to land in between the two locations (Becker and Jürgens 1979). This
74 amplitude transition function can be used to determine the “point of no return” at
75 which the new input can be no longer affect the motor plan and the saccade will not
76 change its destination, and it is defined by the transition point for the first deviations
77 from the first target position (i.e. the earliest sign of any influence of the second
78 target). This time interval was interpreted as the delay between the afferent signal
79 reaching the first oculomotor structures and the triggering of the eye movement signal
80 to the muscles (Becker 1991).

81 The period between this point and saccade onset, which Ludwig et al (2007)
82 called “saccadic dead time” (SDT), has been estimated to be as brief as 60 ms
83 (Findlay and Harris 1984; Ludwig et al. 2007). At first, it was suggested that the SDT
84 was a constant value, about 70 ms, similar across different eye movement tasks
85 (Beutter et al 2003; Findlay and Harris 1984; Hooge et al 1996; Ludwig et al 2005;
86 Van Loon et al 2002) and it also represented a critical parameter for models of eye
87 movements (Nuthmann et al 2010; Reichle et al 1998; Van Loo et al 2002). More
88 recently, this notion was challenged by Ludwig et al (2007) by showing that even if
89 the SDT was not influenced by variations in saccadic reaction times it was susceptible
90 to manipulations of the spatial configuration of the two targets. Similarly, Walshe and
91 Nuthmann (2015) showed that the SDT was affected by the type of background used
92 during double-step tasks, approaching a minimum value of 70 ms for uniform scenes

(black background). Nonetheless, the lower limit reported in behavioural studies has not been lower than the 60 ms estimated by Ludwig et al (2007). The onset of SI, 60-70 ms after a visual flash, is thus compatible with the concept of saccadic dead time, implying a generalized temporal boundary before saccadic execution, during which new visual changes, either relevant (double-step) or irrelevant (distractors), cannot influence the impending saccade.

Across multiple studies of SI, Reingold and Stampe reported consistent changes in the timing of saccades, accounting for the SI dip profile, but they did not report any spatial or kinematic changes in the saccades that were launched. More recently, however, there have been clear indications that SI does have some influence on the spatial aspect of saccadic behavior (Buonocore and McIntosh 2012; Edelman and Xu 2009; Guillaume, 2012). Specifically, saccades launched during the period immediately preceding or following the SI dip, induced by a contralateral distractor or a mask covering a large part of the screen and target, have been found to be hypometric (falling short of the target) (Edelman and Xu 2009; Guillaume, 2012). These observations may echo findings made in studies of micro-saccades (Hafed and Ignashchenkova 2013; Rolfs et al. 2008). For example, Hafed and Ignashchenkova (2013) reported that the micro-saccadic rate was not only reduced 100 ms after a supplementary stimulus, but that the spatial character of the persisting micro-saccades was sensitive to the location of that stimulus. Their interpretation was that the observed micro-saccades reflected an instantaneous “read out” of activations in the oculomotor maps of the superior colliculus, affected both by the target and the supplementary stimulus.

Recent literature thus suggests that SI might not be exclusively temporal in nature but may also involve changes in the kinematic and spatial aspects of the

118 saccade. However, aside from the work of Guillaume (2012) and some observations
 119 made by Edelman and Xu (2009), no other studies have made a detailed analysis of
 120 the time-course of such parameters following distractors with different characteristics,
 121 in a way that is analogous to what has been done for the temporal domain.

122 In the present paper we adopted precisely this strategy. In addition to a
 123 standard SI analysis, we applied a time-course analysis to study the gain and peak
 124 speed of saccades launched at different times following a visual flash. First, we
 125 applied this novel analysis to a previously collected dataset (from an unpublished
 126 experiment that incorporated SI within a visual discrimination task) that was well
 127 suited to this exploration. This exploratory Experiment 1 confirmed that SI could be
 128 associated with a modulation of saccadic gain, following a central flash not dissimilar
 129 to the transient mask that Guillaume (2012) found to affect saccadic amplitude. We
 130 followed up this preliminary observation with two experiments designed to more
 131 finely measure the subtle changes in the saccade characteristics. In Experiment 2, we
 132 used a generalized flash located in the top and bottom of the screen to exclude the
 133 possibility that the amplitude effects were related specifically to the spatially localized
 134 nature of the central flash. In Experiment 3, we manipulated distractor location, to be
 135 more or less eccentric than the target, to test whether saccade hypometria was
 136 dependent upon distractor location, as has been suggested for micro-saccades (Hafed
 137 and Ignashchenkova 2013), or resulted from a more general inhibitory phenomenon.
 138 Across these three experiments, we report a complex interplay between spatial and
 139 temporal modulations for distractors interfering at different stages of saccade
 140 programming and execution, including during the commonly accepted “saccadic dead
 141 time” that is thought to occur after the saccade plan passes a point of no return.

142

143 **Method**

144

145 *Participants*

146 Nine (Experiment 1), ten (Experiment 2) and eight (Experiment 3) volunteers aged
147 between 18 and 30 years participated. All were free from neurological and visual
148 impairments. The experiment was conducted in accordance with the 1964 Declaration
149 of Helsinki, and the guidelines of the University of Trento Research Ethics
150 Committee for behavioral experiments. All participants gave informed written consent
151 and received €7 per testing hour, or course credits.

152

153 *Apparatus, stimuli and procedure*

154 Stimuli were presented on a 17-inch CRT monitor (1024 x 768 pixels) at 85 Hz
155 (Experiment 1) or 100 Hz (Experiment 2 and 3). In all the experiments, participants
156 were seated with their head resting on a chin and forehead rest in order to reduce head
157 movements. The eyes were horizontally and vertically aligned with the center of the
158 screen at a distance of 60 cm. Eye movements were recorded with the EyeLink 1000
159 system (detection algorithm: pupil and corneal reflex; 1000 Hz sampling; saccade
160 detection was based on a 30 deg/s velocity and 9500 deg/s² acceleration thresholds;
161 maximum head movement¹ tolerance equal to 25 mm by 25 mm by 10 mm -
162 horizontal by vertical by depth respectively). In all three experiments, a five point-
163 calibration on the horizontal and vertical axes was run at the beginning of each
164 session and after three consecutive trial blocks. Additional calibrations were added if
165 the participant moved their head from the chinrest. In all the experiments the
166 background was grey (23.5 cd/m²). The experimenter started each trial with a drift

167 correction, after which a tone accompanied the onset of a 0.50° central fixation cross
168 (124 cd/m^2).

169 In *Experiment 1* (Figure 1A), after a random interval varying between 500 to
170 1200 ms, a red dot (0.5° , 28.2 cd/m^2) was displayed at 10° of eccentricity, equally
171 often to the right or to the left of fixation. Participants were required to make a
172 saccade to this target as soon as it appeared. Independent of that requirement, in half
173 of the trials, a black square was flashed for 11.7 ms at the center of the display; this
174 square was the “flash” stimulus used to elicit SI in this experiment (see below).
175 Saccadic reaction times (SRT) were recorded as the interval between target onset and
176 the start of the saccade. This first experiment was originally designed for a different
177 purpose and incorporated a perceptual task whereby four Gabor patches (size = 6° ;
178 frequency = 0.9 cycle/degree) were presented in the four corners of the monitor for 12
179 ms, 105 ms after flash (or invisible flash in target only condition) onset. On half of the
180 trials, the four stimuli had the same orientation (vertical or horizontal) and on the
181 other half, one of them had a different orientation. At the end of the trial, participants
182 were asked to report if all the Gabor patches were the same or if one was different.
183 This perceptual element of Experiment 1 is not relevant for present purposes and the
184 results of the perceptual task were analyzed separately in a different unpublished
185 manuscript focusing on saccadic suppression. Critically, the present analyses were
186 restricted to trials in which saccades were launched up to 45 ms after the display of
187 the Gabor patches. Thus, the presence of the perceptual targets was not likely to
188 influence the pattern of results. No perceptual targets were present in the other two
189 experiments reported here.

190 Participants performed a preliminary block of 64 target-only trials, half with
191 the target on the right and half with the target on the left side of the screen. The

192 median SRT from the last 50 of these trials provided an estimate of the expected SRT
193 for that participant for the experimental blocks. In the experimental blocks, target-
194 only (no-flash) trials were intermingled equally with target plus distractor (flash)
195 trials, in which, in addition to the target, the black square (3.5° , 2.3 cd/m^2) was
196 flashed at the center of the screen for 11.7 ms. The onset of this central flash varied
197 randomly between ~ 117 ms before to 11.7 ms after the expected SRT for that
198 participant in steps of 11.7 ms, thereby providing a wide range of distractor delays.
199 Each of the two conditions (flash, no flash) occurred 64 times per block, shuffled
200 randomly. Each participant completed two sessions of eight experimental blocks, on
201 different days, for a total of 1024 trials. Although originally conceived for a different
202 purpose, the experimental design described above provided a rich dataset for an
203 opportunistic exploration of the time-course analysis of saccade kinematics, and
204 provided the basic template for the two experiments subsequently designed to further
205 investigate these issues (but which did not include the perceptual task, and utilized
206 different distractor locations).

207 Experiment 2 (Figure 1B) was designed to replicate and extend the
208 observations of the first experiment. The saccadic task was similar to that of
209 Experiment 1 but changes were introduced to sample a wide range of distractor delays
210 relative to expected saccade onset and to optimize the effect of the visual transient
211 upon the oculomotor response. No perceptual task was presented in either the second
212 or third experiment. After a random interval varying between 500 to 1200 ms, a white
213 dot (0.5° , 124 cd/m^2) was displayed at three possible eccentricities (4° , 8° and 12°
214 degrees of visual angle) with equal probability to the right or to the left of the visual
215 field. As in Experiment 1, each participant performed a preliminary block of 60
216 target-only trials (20 for each eccentricity) to determine the median SRT from which

217 to calculate flash onset. In the experimental blocks, target-only (no-flash) trials were
 218 intermingled with target plus distractor (flash) trials. The distractor consisted of two
 219 white rectangles (width: $\sim 33^\circ$, length: $\sim 8.5^\circ$, 124 cd/m^2) covering one third of the top
 220 and bottom of the screen (see: Reingold and Stampe (2002) for a similar procedure).
 221 The flash was presented for 20 ms. During the course of the experiment, flash onset
 222 was varied around the participant-specific median SRT by randomly subtracting one
 223 of six possible SOAs, spanning from 20 to 120 ms in steps of 20 ms. In each trial we
 224 recorded the SRT and, at the end of the trial, calculated flash-to-saccade delay for that
 225 trial by subtracting flash onset from the current SRT. To ensure adequate sampling of
 226 saccades in each time bin after flash onset (bin size 20 ms), we kept track of the
 227 number of saccades recorded within each time bin and, when any bin reached a
 228 threshold of 60 observations, replaced the SOA most closely matching that flash-to-
 229 saccade delay with the SOA of the least represented bin. At the end of each block, the
 230 median SRT, used to calculate flash onset, was updated with the median of the current
 231 block. Overall, we ran 260 trials per condition, i.e. two flash (absent-present)
 232 conditions by three target eccentricities (4° , 8° and 12° of visual angle) for a total of
 233 1560 trials. Participants completed two sessions on different days in which the 780
 234 trials were divided in 13 blocks of 60 trials each.

235 Experiment 3 had a similar procedure to Experiment 2 but only one target
 236 eccentricity was used (10° of visual angle) (Figure 1C). As in the other experiments,
 237 each participant performed a preliminary block of 30 target-only trials to determine
 238 the median SRT from which to calculate flash onset. In the experimental blocks,
 239 target-only (no-flash) trials were intermingled with target plus distractor (flash) trials.
 240 In distractor trials, the flash was presented for 20 ms and consisted of a white vertical
 241 rectangle (width: $\sim 2^\circ$, length: $\sim 24^\circ$, 124 cd/m^2) either less eccentric (flash-, 6°) or

242 more eccentric (flash+, 14°) than the target (10°). Flash onset was varied around the
243 per-participant median SRT, which was updated after each block, by randomly
244 subtracting one of four possible SOAs spanning from 30 to 60 ms in steps of 10 ms.
245 This procedure generated a high density distribution within the first 130 ms after flash
246 onset allowing us to strengthen the analysis of amplitude and velocity variations.
247 Overall, we ran 20 trials per flash condition (flash absent, flash- and flash+) for a total
248 of 60 trials per block. Participants completed two sessions for eight blocks in one day
249 for a total of 960 trials, with 320 trials per flash condition.

250

251 *Data screening*

252 We excluded saccades with latencies of less than 70 ms (Experiment 1: ~1.2%;
253 Experiment 2: ~2.5%; Experiment 3: ~0.73%) or of more than 500 ms (Experiment 1:
254 ~2.7%; Experiment 2: ~0.14%; Experiment 3: ~0.75%). We also removed saccades
255 with an amplitude less than 1° amplitude (Experiment 1: ~0.6%; Experiment 2: 3.4%;
256 Experiment 3: ~2.10%) and saccades made in the wrong direction (Experiment 1:
257 ~0.02%; Experiment 2: ~2.4%; Experiment 3: ~0.05%). In Experiment 2 and 3 we
258 also excluded 2.6% and 1.52% of saccades, respectively, due to blinks.

259

260 *Analysis of saccadic inhibition*

261 In all the Experiments, we performed an analysis of the SRT distributions for all valid
262 trials by following Bompas and Sumner's (2011) procedure to calculate the "dip"
263 ratio. As a first step, we recoded SRTs relative to flash onset, by subtracting from
264 each SRT the SOA between target and flash. Then, for each participant and condition
265 (Experiment 1: no-flash and flash; Experiment 2 no-flash and flash at each target
266 eccentricity; Experiment 3: no-flash, flash- and flash+), we created percentage

frequency histograms (bin width 4 ms) that were then lightly smoothed using a Gaussian kernel with 24 ms window and 2 ms SD. The smoothed histograms were interpolated to obtain 1 ms precision. To estimate the level of SI, we computed the proportional change for each point in time in the flash distribution relative to the no-flash distribution by using the formula: $(\text{no-flash} - \text{flash})/\text{no-flash}$. This operation was performed on both the no-flash and flash condition. In the no-flash condition an “invisible” stimulus was presented using the same time procedure as for the flash condition (for detailed analysis on this procedure see: McIntosh and Buonocore 2014). The magnitude (i.e. maximum of inhibition) and the latency (time to the maximum) of SI were taken in the first 150 ms after flash onset. To visualize the average SI profile, per condition, the individual profiles were then averaged across participants and the 95% confidence interval was computed at each time point (Figure 3A, D and G). Statistical analysis was performed on the individual parameters extracted from each SI profile across the three Experiments are reported in Table 1, 2 and 3.

Analysis of saccadic kinematics

The analysis of saccade kinematics focused upon saccadic gain and normalized peak speed. The first step was to extract these variables for every trial. Saccadic gain is saccade amplitude divided by the target amplitude for that trial, with values greater than one indicating overshoot (hypermetria), and values less than one indicating hypometria (undershoot). Normalized peak speed was the *observed* peak speed divided by the peak speed *predicted* from the observed saccade amplitude on that trial, with values bigger than one indicating a speed higher than expected, and values less than one indicating a speed lower than expected. The calculation of normalized

292 peak speed therefore included an additional initial step to predict peak speed from the
 293 main sequence relationship between saccade amplitude and peak speed. To do so, for
 294 each participant separately, we fitted a polynomial function to the *observed* peak
 295 speed over the *observed* saccadic amplitude in all no-flash trials, and extracted the
 296 polynomial for the best fit according to a least-squares procedure. In Experiment 1
 297 and Experiment 3, there was only one target location, and the spread of observed
 298 saccade amplitudes was too small ($\sim 2^\circ$) to model the entire main sequence function,
 299 so we used a 1st order polynomial function. In Experiment 2, we made use of all the
 300 eccentricities to estimate the best main sequence fit using a 2nd order polynomial
 301 function. In Figure 2 we show one example of fitting for each experiment (panel A, B
 302 and C) along with the R^2 for each participant in all of the experiments. Based on these
 303 individual fit parameters, we derived the *predicted* peak speed from the observed
 304 saccade amplitude in each trial, and used this value to normalize the observed peak
 305 speed for that trial.

306 We then analyzed the time-course of these kinematic variables relative to the
 307 flash event inducing SI. For each participant, RTs were binned using a bin-width of
 308 20 ms and the mean saccadic gain and normalized peak speed was calculated for
 309 saccades launched within each time bin. For Experiments 1 and 3, the means were
 310 entered into separate two (flash: no-flash vs. flash) by seven (bin: 10 to 130 in 20 ms
 311 intervals) repeated-measures ANOVAs. In Experiment 2, a two (flash: no-flash vs.
 312 flash) by three (eccentricity: 4° , 8° and 12°) by seven (bin: 10 to 130 in 20 ms
 313 intervals) repeated-measures ANOVA was performed, with Greenhouse-Geisser
 314 adjustments to the degrees of freedom where sphericity was violated. Significant
 315 interactions were followed up by a series of paired samples t-tests comparing no-flash
 316 versus flash conditions at each time bin. Considering that adjacent time bins are likely

to be correlated, we performed the Benjamini & Hochberg (1995) and the Benjamini & Yekutieli (2001) procedure controlling the false discovery rate (FDR) of a family of hypothesis tests. Corrected p-levels are reported in the text.

Results

Experiment 1 - Analysis of saccadic inhibition

Overall, we confirmed the main SI effect by showing a strong bimodality in the flash histogram, with the lowest saccadic frequency happening around 90 ms after flash onset. For illustrative purposes, Figure 3A shows the average SI profile across participants, expressed as the ratio of inhibited saccades (i.e. delayed) to baseline saccadic frequency for the no-flash condition (see Methods section for details of the SI profile calculation). Using the parameters extracted from the individual SI profiles, we estimated that an average maximum of 78% of saccades were inhibited at 85 ms after the flash onset, matching well with the timing of SI and micro-saccade inhibition found in previous experiments (Bompas and Sumner 2011; Buonocore and McIntosh 2008, 2012, 2013; Edelman and Xu 2009; Guillaume et al. 2012; Hafed and Ignashchenkova 2013; Reingold and Stampe 2002). Individual values for the latency and the magnitude of inhibition were consistent across participants (Table 1).

Experiment 1 - Analysis of saccadic kinematics

For the gain, the repeated-measures ANOVA revealed a main effect of Flash [$F(1,8) = 12.96; p < 0.01$] and Bin [$F(2.25,17.99) = 8.29; p < 0.005$] but more interestingly, there was a significant interaction between the two factors [$F(1.6,12.82) = 8.99; p < 0.005$] (Figure 3B). During flash trials we observed a strong decrease in saccadic

342 amplitude (hypometria) for saccades launched 20 to 80 ms after flash onset (30 ms
 343 bin: [$t(8) = 6.29$; $p < 0.002$]; 50 ms bin [$t(8) = 3.92$; $p < 0.016$]; 70 ms bin [$t(8) =$
 344 3.35 ; $p < 0.023$]). In order to estimate a possible violation of the main sequence, we
 345 analyzed the time-course of the normalized peak speed. We report a significant main
 346 effect of Flash [$F(1,8) = 10.00$; $p < 0.01$] and Bin [$F(6,48) = 4.03$; $p < 0.005$] but no
 347 interaction between these factors [$F(3.22,25.78) = 2.21$; $p = 0.1$]. The data suggest a
 348 general disturbance of the main sequence during flash trials, with peak speed
 349 exceeding the value predicted from saccadic amplitude. Looking at Figure 3C, there is
 350 an indication that the violation might be concentrated in a few specific time-points
 351 after flash onset, during the pre- and post- inhibitory period, as observed by
 352 Guillaume (2012). Nonetheless, while these data are suggestive, we were unable to
 353 confirm a significant temporal modulation of the main sequence. However, it should
 354 be noted that the above was an opportunistic and exploratory analysis of a dataset
 355 collected for different reasons. Experiments 2 and 3 directly investigated these trends
 356 with more targeted studies that were designed to have greater power to investigate the
 357 kinematic changes suggested by Experiment 1.

358

359 *Experiment 1 - Interim discussion*

360 Taken together, the modulations in saccadic gain and normalized peak speed suggest
 361 a general violation of the main sequence. First, a strong saccadic hypometria was
 362 observed for saccades launched between 30 (-0.48°) and 70 ms (-1.20°) after flash
 363 onset, which was not accompanied by a proportional reduction in peak speed. This
 364 pattern of data suggests that saccades launched during this period may have initially
 365 been programmed for greater amplitudes, but terminated in-flight following arrival of
 366 the distractor signal (Edelman and Xu 2009; Guillaume 2012; Munoz et al. 1996).

367 There were subsequent, albeit weaker, indications of a second violation toward the
 368 end of SI, where the peak velocity tended to exceed that predicted from the main
 369 sequence. Considering that the average no-flash SRTs were ~220 ms for this task,
 370 saccades launched 130-150 ms after flash onset corresponded to flash stimuli
 371 presented only 70 - 90 ms after target onset, thus relatively close in time to the target
 372 onset. The presentation of the flash may thus have summed with the build-up of
 373 target-related activity, generating an overall increase in the level of SC activation. At
 374 the time of saccade launching, this increased activity might have translated as
 375 increased velocity.

376 The exploratory analysis reported above brought up an intriguing pattern of
 377 modulations following distractor interference that confirmed and expanded previous
 378 reports of spatial and temporal effects (Edelman and Xu 2009; Buonocore and
 379 McIntosh 2012; Guillaume (2012). Nonetheless, while reduced saccadic gain was
 380 clear during the pre-inhibitory period (Edelman and Xu 2009; Guillaume 2012), the
 381 pattern of elevation of normalized peak speed was not so tightly locked to a particular
 382 time period; a more powerful experiment may be required to determine these patterns
 383 of kinematic variation more definitively. Moreover, in the present experiment we used
 384 as the distractor a single, highly localized and central flash that might have interfered
 385 with saccadic amplitude during target selection because it was partially interfering
 386 with the saccade trajectory, similarly to the mask stimuli used by Guillaume (2012).
 387 Instead of causing general inhibition, this less eccentric distractor might have induced
 388 smaller saccadic amplitudes via spatial interference, offering an alternative account of
 389 the observed hypometria. This could be analogous to observations of micro-saccadic
 390 inhibition, whereby the target-flash configuration was found to determine the pattern
 391 of amplitudes changes (Hafed and Ignashchenkova 2013; Rolfs et al. 2008).

Thus, to more closely measure the possible violations of the main sequence found in this preliminary dataset, we designed a further experiment to test whether these patterns were robust. First, we increased the power to detect small variations by substantially increasing the number of trials. Second, the timing of the flash was more finely tuned online to each participant's saccadic performance in order to elicit a strong SI in every participant. Third, to minimize the possibility of a direct spatial interference of the distractor as a competing saccadic target, the flash was more spatially generalized across the display, occupying both the top and bottom thirds of the screen (see Reingold and Stampe, 2002). Finally, we extended the range of target eccentricities to better map the main sequence function.

402

403 *Experiment 2 - Analysis of saccadic inhibition*

The parameters extracted from the individual SI profiles (Table 2) were closely similar across the three eccentricities and the maximum inhibition was about 74, 74 and 77 percent for the three eccentricities respectively with a latency of 78, 79 and 77 ms after the flash onset, matching the data from Experiment 1. Neither the magnitude nor the latency of inhibition were significantly different between the three eccentricities [magnitude: $F(2, 18) < 1$; N.S.; latency: $F(1.13, 10.15) < 1$; N.S]. For descriptive purposes, in Figure 3D we report the average profile across the three eccentricities.

412

413 *Experiment 2 - Analysis of saccadic kinematics*

By using the gain as measure of saccadic spatial performance, we found a significant main effect of Flash [$F(1,9) = 13.13$; $p < 0.006$] and Bin [$F(1.943, 17.488) = 5.64$; $p < 0.01$] and again a significant interaction between the two factors [$F(2.361, 21.245) =$

5.6; $p < 0.008$] (Figure 3E). There was no effect of eccentricity, suggesting that these modulations were similar across a range of saccadic amplitudes. The gain was reduced for saccades launched at 30 ms after flash onset [$t(9) = 3.41$; $p < 0.027$] with a minimum value for saccades launched at 50 ms after flash onset [$t(9) = 5.39$; $p < 0.003$], replicating the finding of Experiment 1. To check if these modulations violated the main sequence, we inspected the normalized peak speed. We report a significant interaction between Flash and Bin [$F(6,54) = 4.65$; $p < 0.001$] (Figure 3F). Pair-wise t-test comparisons confirmed a violation exceeding the expected peak speed for saccades launched at 30 ms [$t(9) = 3.41$; $p < 0.0273$]. More anomalously, there was a significant *reduction* in normalized peak speed for saccades launched at 50 ms after the distractor [$t(9) = 4.28$; $p < 0.014$]. Overall, the data from Experiment 2 confirmed and extended the results reported in Experiment 1. We replicated saccadic hypometria during the pre-inhibitory period (Edelman and Xu 2009; Guillaume 2012), associated with a violation of the main sequence and confirmed that this main sequence violation was specific in time. We again saw a qualitative trend toward a second, later rise in the main sequence ratio during the post-inhibitory period, although this trend did not reach statistical significance.

434

435 *Experiment 2 - Interim discussion*

436 The data from Experiment 2 confirmed that saccades launched during the pre-
 437 inhibitory period were truncated in flight, perturbing the main sequence (Edelman and
 438 Xu 2009; Guillaume 2012). Additionally, and surprisingly, we also observed a
 439 reduction of the normalized peak speed just before the start of inhibition. This finding
 440 was unexpected and, at present, we do not have a firm explanation for it. One
 441 possibility is that, on entering into the inhibitory period, when the interference is

442 maximal and the reduction in gain is peaking, saccades may be truncated even *before*
443 achieving peak speed, consequently decreasing the ratio between the predicted and the
444 observed velocities. This would predict that saccades launched in this time period
445 would be associated with a reduced duration, since the truncation would happen so
446 early. To explore this idea, we ran an analysis of saccadic duration, and confirmed a
447 significant reduction specifically for saccades launched at 50 ms after the distractor
448 [$t(9) = 3.58$; $p < 0.0417$], thus coincident with the reduced peak speed. Nonetheless,
449 since this pattern of reduced peak speed was not evident in Experiment 1, more
450 studies are needed to rule out the possibility that this observation was just a chance
451 finding. Finally, we again saw indications, albeit relatively weak, of violations of the
452 main sequence during the post-inhibitory period.

453 In Experiment 1, we considered that one possible explanation for the reduction
454 in saccadic gain was that the flash-related activation may have interfered directly with
455 the planning of the saccade trajectory; that is, a spatial averaging effect. In
456 Experiment 2, this issue was addressed by placing the flash in the top and bottom
457 third of the screen (Reingold and Stampe, 2002). Nonetheless, one could argue that
458 the “center of gravity” of the flash configuration was still at the center of the screen;
459 according to the micro-saccade inhibition literature, the final read out of the superior
460 colliculus activation after flash presentation could be skewed toward the screen
461 center, predicting hypometria by spatial averaging.

462 To better test the possibility of a spatial averaging effect we ran Experiment 3
463 in which the position of the flash relative to the target was either less (flash-) or more
464 (flash+) eccentric than the saccade target. If the hypometria was generated by a
465 general truncation mechanism, we should see the hypometria for both the less and
466 more eccentric flash. On the other hand, if the effect is driven by flash location we

467 should record hypometria for the less eccentric flash and hypermetria for the more
468 eccentric flash.

469 Although Edelman and Xu (2009) tested the effect of distractor location on SI,
470 reporting that flashes appearing at the location of the saccade goal led to “express-
471 like” saccades, rather than SI, no prior study (cf. Guillaume, 2012) has systematically
472 investigated the effect of the flash location relative to saccadic target upon saccadic
473 amplitude and peak speed, leaving this important issue open.

474

475 *Experiment 3 - Analysis of saccadic inhibition*

476 The SI profile in the flash+ condition was smaller compared to the flash- condition
477 (Figure 3G). The analysis performed on the parameters extracted from the individual
478 profiles showed that the maximum inhibition was about 52% (flash+) and 83% (flash-
479) [$t(7) = 6.71$; $p < 0.0005$] with a latency of 74 and 71 ms respectively after the flash
480 onset [$t(7) = 1.26$; N.S.]. Individual parameters for the two conditions are reported in
481 Table 3. These data imply that the eccentricity of the flash, relative to the target, has a
482 strong impact on the level of saccadic inhibition, an interesting observation that has
483 been little explored in prior studies.

484

485 *Experiment 3 - Analysis of saccadic kinematics*

486 For gain, the main effect of Flash was reliable [$F(2,14) = 47.92$; $p < 0.0001$] as was
487 the main effect of Bin [$F(6,42) = 5.12$; $p < 0.001$]. More importantly, as in
488 Experiments 1 and 2, there was a significant interaction between the two factors
489 [$F(12,84) = 16.67$; $p < 0.0001$] (Figure 3H). Follow-up analyses of the Flash by Bin
490 interaction replicated the strong hypometria effect but with different timings for the
491 two conditions. In the flash- condition, the hypometria started for saccades launched

30 ms after flash onset and numerically peaked for saccades launched at 50 and 70 ms after flash onset [Bin 30: $t(7) = 5.30$; $p < 0.0039$; Bin 50: $t(7) = 7.60$; $p < 0.0009$; Bin 70: $t(7) = 4.19$; $p < 0.0095$] (as for Experiment 1 and 2). On the other hand, hypometria was observed, but started much later in the flash+ condition, peaking for saccades launched between 110 to 130 ms after flash onset [Bin 110: $t(7) = 7.1178$; $p < 0.0013$; Bin 130: $t(7) = 5.5058$; $p < 0.0032$]. These pronounced differences of timing allow the possibility that the two types of hypometria might have different origins.

As for the other experiments, the reductions in gain were accompanied by violations of the main sequence. We report a significant main effect of Bin [$F(6,42) = 7.63$; $p < 0.001$] and a significant interaction between Flash and Bin [$F(12,84) = 5.51$; $p < 0.0001$] (Figure 3I). In particular, the violation was present for saccades launched at 30 ms in the Flash- condition [$t(7) = 3.92$; $p < 0.020$] and followed by violations in the post-inhibitory period at 110 ms and 130 ms [$t(7) = 3.03$; $p < 0.044$, $t(7) = 5.79$; $p < 0.005$]. The flash+ condition had only one significant violation point during the post-inhibitory period, at 130 ms [$t(7) = 3.9773$; $p < 0.0374$]. We did not see in any of the conditions a reverse in the violation, as observed in Experiment 2. Nonetheless, looking at the bottom row of Figure 3 it is suggestive that for all the experiments the shape of the normalized peak speed oscillated compared to the steady baseline condition, with higher or lower values alternating within the total time-course. Thus, although the most consistent statistical pattern is for distractor-induced reductions in saccadic gain, with violations of the main sequence in a positive direction (i.e. increased peak speed to amplitude ratios), the qualitative pattern emphasizes that the perturbations of the main sequence may be somewhat unstable in direction as well as degree.

Overall, this pattern of results suggests that the variations in amplitude might be first driven by a truncation mechanism followed by a readout of the superior colliculus map, similarly to what has been reported in the micro-saccade literature (Hafed and Ignashchenkova 2013). The flash- condition showed a clear truncation (hypometria accompanied by relatively high peak speed) stopping the saccade in-flight for motor programs launched 30 ms after flash onset. This was followed up by a strong hypometria (but with appropriately-scaled peak speed), as predicted by saccadic averaging. On the other hand, in the flash+ condition there was no significant evidence of hypometria or increase in peak speed soon after flash onset. Moreover, the kinematics of saccades launched during the SI period were not influenced by the presence of the flash. The very large difference between the two flash conditions indicates that the spatial layout was having an impact on saccadic amplitude in a way compatible to a spatial readout of the superior colliculus map. Nonetheless, contrary to a strict prediction of the read out hypothesis, we do not report any hypermetria for the flash+ condition, but this was probably a simple consequence of the logarithmic compression of the visual map in which more eccentric locations occupy less neural tissue (Ottes et al. 1986; Van Gisbergen et al. 1987).

A final interesting observation is that we also recorded hypometric saccades in the flash+ condition, but *following* the inhibitory period. The hypometria was also accompanied by an increased peak speed indicating that these saccades were programmed for the correct target location but subsequently felt short, leading to a violation of the main sequence. It is important to note that these saccades were ones that would have been re-instated or reprogrammed, so the reported effect is not the same as the hypometric saccades recorded during the pre-inhibitory period. This late hypometria is more similar to the one reported by Guillaume (2012) with masking

stimuli covering either the entire screen (full mask) or only the portion of the screen where the target was displayed (half mask). Similarly to our findings, Guillaume also observed an increase in peak speed for these reinstated saccades, as in our Experiment 1, 2 and 3 (where we did not record a gain reduction). One possibility might be that the later spatial effects are generated by cortico-tectal feedback from areas such as the frontal eye field and the lateral intraparietal cortex inhibiting the SC and truncating the saccade at a later processing stage.

General conclusions

In three experiments, we flashed a visual transient at a range of times relative to a target-directed saccade, at different positions: either at fixation, at the top and bottom of the screen or at a location on the target axis more or less eccentric than the target. In all cases, once the data were aligned temporally to the onset of the flash, a distinctive pattern of variation in saccadic behavior was revealed both in *time* and *space*.

First, we replicated the well-known temporal inhibitory effect of the flash (SI: Reingold & Stampe, 1999, 2002) on the initiation of saccades, with a maximal decrease in saccadic frequency varying from 53 to 83 percent across experiments, and the latency of maximum inhibition ranging from 77 to 86 ms. The decrease in saccadic frequency began as early as 60 ms, recovering by 110 ms after the flash. These timings are compatible with the idea that the triggering mechanism of a saccade cannot be changed beyond a point-of-no-return around 60 ms before launching (Reingold and Stampe 2002), and thus with the concept of a “saccadic dead time” applied to this pre-launch period (Findlay and Harris 1984; Ludwig et al. 2007). In passing, we also made a new observation (Experiment 3) that SI magnitude, but not

latency, was strongly affected by flash eccentricity, with greater inhibition for nearby distractor locations. This result, although not a focus of our paper, carries the interesting suggestion that eccentricity is more influential upon SI than distance from target (since in our experiment the flash was equally distant from the target in both the flash+ and flash- conditions).

Our major interest was in the kinematic character of saccades launched following a flashed distractor, and here we focused on saccade amplitude (gain) and its relation with peak speed (main sequence relation). In all Experiments, we observed a strong hypometria for saccades launched a mere 20 ms after the flash, extending to saccades launched up to 80 ms after the flash. The maximum reduction in gain was ~12% in Experiment 1, ~5% (considering all target eccentricities together) in Experiment 2, and ~15% in Experiment 3. Interestingly, the hypometric saccades were not always accompanied by the correspondingly lower peak speed expected from the main sequence. These perturbations of the main sequence were time specific in both Experiments 2 and 3, and maximal for saccades launched around 30 ms after flash onset. A second peak of relative increase in the peak speed was visible for saccades launched around 110 and 130 ms after the flash, in this case unaccompanied by an increase in saccadic gain. Taken together, the data show a complex violation of the main sequence around the onset and offset of the SI dip that develops over time, oscillating with higher or lower values compared to the steady baseline condition.

One hypothesis to account for the early perturbation of the main sequence during the pre-inhibitory period (reduced gain without reduced peak speed) would be that a saccade already in flight was suddenly interrupted by flash onset (see also: Edelman and Xu 2009; Guillaume 2012), creating hypometric saccades with peak speeds appropriate to the originally intended target. In Experiment 2, we additionally

592 observed a decrease of the normalized peak speed compared to baseline toward the
593 end of this early period of perturbation, suggesting that in some circumstances the
594 saccades might have been truncated prior to achieving the peak speed expected for
595 that amplitude. The most striking aspect of these data is saccadic modulation for
596 distractors presented a mere 30 ms before execution, and thus 30 ms before the
597 earliest inhibition of saccade launching. This demonstrates interference from
598 distractors presented during “saccadic dead time” (Ludwig et al. 2007; Weber et al.
599 1992), in which the saccadic program is past the “point of no return” (Reingold &
600 Stampe, 2002), and should be impervious to further visual stimulation. It may indeed
601 be that no changes were implemented to the saccade program itself, but that these
602 very late distractors may have acted to modify the saccade in-flight. Our result
603 confirms that this terminal phase of saccade preparation, immediately prior to
604 launching, despite being immune to reprogramming, may still be permeable to
605 distractor interference during saccade execution, beyond the point of no return.

606 The late phase of kinematic perturbation, around the offset of the SI dip, had a
607 rather different character. We found a pattern of elevated peak speeds without a
608 significant change in saccadic amplitude, except for the flash+ condition in
609 Experiment 3. This late phase of perturbation was visible in all three experiments, but
610 was statistically weak, reaching significance only in Experiment 3. One speculation is
611 that this reflects something about saccades being recovered, or reprogrammed
612 following inhibition, as if these inhibited saccades required an additional impetus to
613 escape the inhibitory effect that resulted in a higher peak speed. Alternatively, the
614 presentation of the flash, temporally close to the target onset for this time period,
615 might have summed up with the target activity leading to an increase in the level of
616 SC activation. These saccades might have remained spatially accurate rather than

617 being hypermetric because of the feedback loop that controls the saccades within the
 618 brainstem (Sparks 2002). Saccades can maintain amplitude information and vary
 619 duration/velocity to compensate for external perturbation, such as in the interrupted
 620 saccades paradigm (Keller and Edelman, 1994). One hypothesis could be that about
 621 100 ms after flash onset the processing of saccadic amplitude was well advanced so
 622 that amplitude/direction were already specified by the activity at the saccadic goal
 623 (Anderson, Keller, Gandhi and Das 1998). Nonetheless, the sudden activation of other
 624 superior colliculus neurons summed up with the ongoing process, resulting in a
 625 “global higher activity” at the time of saccade launching, that we recorded as
 626 increased velocity, as predicted by the “dual coding” (Sparks and Mays 1990) and
 627 vector summation hypotheses (Goossens and Van Opstal 2006; Van Opstal and
 628 Goossens 2008). Further replication work and modeling of the activity within the
 629 superior colliculus layers would be required before advancing any strong functional
 630 interpretation of this late perturbation of the main sequence.

631 Alternatively, according to Guillaume (2012) the second modulations are
 632 related to the activity induced by the flash in cortical areas sending inhibitory signal to
 633 the SC. This second mechanism would interrupt saccades similarly to the early
 634 mechanisms, hence generating the modulations observed also in the kinematics. Our
 635 data do not fully support the view that the late modulations mimic the truncation
 636 mechanism observed soon after flash onset, as in Guillaume 2012, since aside from
 637 the flash+ condition in Experiment 3, we did not record late hypometric saccades. On
 638 the other hand, we do agree that cortico-tectal feedback, especially from the frontal
 639 eye field, might modulate the motor program during the post-inhibitory period and
 640 have an impact on the spatial parameters of the saccade.

641 In terms of neurophysiology, given its wide generality across tasks and its
642 short latency, SI has been conceptualized as a low-level interference in the early
643 stages of visual processing and it has been modeled in terms of activity within the
644 intermediate and deep layers of the superior colliculus (Bompas and Sumner 2011).
645 Target and flash onsets generate a burst of activation in the superior colliculus
646 oculomotor map. Following the burst, buildup neurons coding for spatially separated
647 target/flash locations (e.g., Everling et al. 1999; Dorris et al. 1997; Munoz and Wurtz
648 1995a) start interacting through lateral inhibition (Olivier et al. 1999). If the flash is
649 central, or not too eccentric, additional stimulation from fixation neurons and/or direct
650 activation of the omnipause neurons might strongly interfere with the completion of
651 the motor program (Gandhi and Keller 1997). In order for a saccade to be inhibited,
652 flash-related interference must begin prior to the “point-of-no-return” at which the
653 saccade-related motor burst is unstoppable (Reingold and Stampe 2002). The latest
654 point in time that a distractor onset can still inhibit saccade execution is determined by
655 the time necessary for visual information to reach the intermediate superior colliculus
656 and to influence motor structures, estimated around 35-47 ms after visual stimulation
657 (Rizzolatti et al. 1980). This timing closely matches the first variation in saccade
658 kinematics, affecting saccades launched around 30 ms after flash onset. Accordingly,
659 in a number of neurophysiological studies with single cell recording from the nucleus
660 raphe interpositus it has been reported that omnipause neurons respond to a light pulse
661 as they do to electrical stimulation, stopping the saccade in flight (Evinger et al 1982).
662 We propose that the early phase of hypometria recorded in the present experiment
663 might have been induced mainly by the sudden activation of the omnipause network
664 subsequent to flash presentation. Another possibility would be that the sudden visual
665 burst elicited by the irrelevant flash interferes with saccade programming to the point

666 that activity for the flash suddenly reaches threshold, favoring interruption of the
667 current saccadic plan, similarly to the mechanisms that generates express saccades
668 (Edelman and Keller 1996).

669 On the other hand, when the transient is presented between 60 to 130 ms
670 before the start of the saccade, the consequences would be expected to be mainly
671 temporal, with a high percentage of inhibited saccades, and the reported hypometria
672 during this phase may reflect the spatial read-out of the SC map. These long lasting
673 inhibitory processes might be driven mainly by lateral inhibition (Buonocore and
674 McIntosh 2008; Olivier et al. 1999; Reingold and Stampe 2002) and reflect
675 competition during target selection processes rather than a sudden truncation of the
676 motor plan.

677 An alternative view, inspired by the micro-saccadic literature, would instead
678 suggest that distractor onset might induce a phase reset. One mechanism that has been
679 proposed to account for the reduction in micro-saccade generation is that the new
680 visual information could generate a countermanding process, cancelling the upcoming
681 micro-saccade in order to initiate a new one (Hafed and Ignashchenkova 2013).
682 Similar processes have been documented for standard saccades within the superior
683 colliculus (Parè and Hanes 2003) and are also compatible with the timings estimated
684 by modeling of SI using competing motor commands (Bompas and Sumner 2011;
685 Trappenberg et al 2001). The stimulus configuration would skew the superior
686 colliculus activity so that saccades would follow the final readout of the superior
687 colliculus activity, predicting modulations in the kinematic parameters similar to
688 those reported here. From the data at hand, we favor the hypothesis that the early
689 hypometria was the consequence of a more general mechanism probably involving
690 the sudden onset of the omnipause neurons network or the activation of burst neurons.

691 Finally, reprogrammed saccades that are launched in the post inhibitory period might
692 have been influenced by extra excitation of the SC map induced by the flash that was
693 temporally close to the target onset.

694 We conclude that distractor effects have broader influences than previously
695 recognized, which can be expressed both in time and space depending on the stage of
696 saccade preparation or execution with which the distractor interferes. Spatial and
697 kinematic effects arise earlier than outright inhibition of the saccade, whilst, more
698 speculatively, saccades reprogrammed after inhibition might exhibit subtly altered
699 kinematics, characterized by increased speed. The point-of-no-return does not put a
700 saccade beyond the reach of distractors; saccades that escape temporal inhibition may
701 instead show changes in velocity, amplitude or both.

702 **Footnotes**

703

704 1. An anonymous reviewer raised the issue of whether the amplitude effects that we
705 observe in our results could be artefactual to head movements. It is important to note
706 that eye movements of this magnitude (lower than 20° of visual angle) are normally
707 accomplished without head movement even in a head un-restrained setup (see
708 Freedman 2008; Fuller 1992). Moreover, it would be very hard to come to a
709 principled account for the exact pattern of gain modulation found here, in particular in
710 the target plus flash conditions, as artefacts of lateralized head movements specific to
711 our experimental manipulation. In Experiment 1 and 2, the stimuli were presented
712 either at the center or top/bottom of the screen, where no lateralized response was
713 required to the distractor.

714 **Acknowledgments.** The research was supported by a European Research
715 Council (ERC) grant (grant agreement no. 313658) awarded to David Melcher and by
716 a research fellowship awarded to Antimo Buonocore and funded by the Comune di
717 Rovereto.
718

719 **References**

- 720 Anderson RW, Keller EL, Gandhi NJ, Das S. Two-dimensional saccade-related
721 population activity in superior colliculus in monkey. *J Neurophysiol* 80:798–
722 817, 1998.
- 723 Bahill A, Clark M, Stark L. The main sequence, a tool for studying human eye
724 movements. *Math Biosci* 24: 191–204, 1975.
- 725 Becker W. Saccades. In: *Vision and Visual Dysfunction*, edited by Carpenter RHS.
726 Boca Raton, FL: CRC, 1991, p. 95–137.
- 727 Becker W, Jürgens, R. An analysis of the saccadic system by means of double step
728 stimuli. *Vision Res* 19(9): 967–83, 1979.
- 729 Benjamini, Y, Hochberg, Y. Controlling the false discovery rate: a practical and
730 powerful approach to multiple testing. *J Roy Statist Soc Ser B* 57: 289–300,
731 1995.
- 732 Benjamini, Y, Yekutieli, D. The control of the false discovery rate in multiple testing
733 under dependency. *Ann Stat* 29: 1165–1188, 2001.
- 734 Beutter BR, Eckstein MP, Stone LS. Saccadic and perceptual performance in visual
735 search tasks. I. Contrast detection and discrimination. *J Opt Soc Am A Opt*
736 *Image Sci Vision* 20: 1341–1355, 2003.
- 737 Bompas A, Sumner, P. Saccadic inhibition reveals the timing of automatic and
738 voluntary signals in the human brain. *J Neurosci* 31: 12501–12512, 2011.
- 739 Buonocore A, McIntosh, DR. Saccadic inhibition underlies the remote distractor
740 effect. *Exp Brain Res* 191: 117–122, 2008.
- 741 Buonocore A, McIntosh, DR. Modulation of saccadic inhibition by distractor size and
742 location. *Vision Res* 69: 32–41, 2012.

- 743 Buonocore A, McIntosh DR. Attention modulates saccadic inhibition magnitude. *Q J*
 744 *Exp Psychol (Hove)* 66: 1051-9, 2013.
- 745 Collewyn H, Erkelens CJ, Steinman, RM. Binocular co-ordination of human vertical
 746 saccadic eye movements. *J Physiol* 404: 183-197, 1988.
- 747 Dorris MC, Pare M, Munoz DP. Neuronal activity in monkey superior colliculus
 748 related to the initiation of saccadic eye movements. *J Neurosci* 17: 8566–8579,
 749 1997.
- 750 Edelman, JA, Keller, EL. Activity of visuomotor burst neurons in the superior
 751 colliculus accompanying express saccades. *J Neurophysiol* 76(2): 908–926,
 752 1996.
- 753 Edelman JA, Xu KZ. Inhibition of voluntary saccadic eye movement commands by
 754 abrupt visual onsets. *J Neurophysiol* 101: 1222–1234, 2009.
- 755 Engbert, R, Kliegl, R. Microsaccades uncover the orientation of covert attention. *Vis*
 756 *Res* 43(9): 1035–1045, 2003.
- 757 Everling S, Dorris MC, Klein RM, Munoz DP. Role of primate superior colliculus in
 758 preparation and execution of anti-saccades and pro-saccades. *J Neurosci* 19(7):
 759 2740–54, 1999.
- 760 Evinger C, Kaneko CRS, Fuchs AF. Activity of omnipause neurons in alert cats
 761 during saccadic eye movements and visual stimuli. *J Neurophysiol* 47: 827–
 762 844, 1982.
- 763 Findlay JM, Harris LR. Small saccades to double-stepped targets moving in two
 764 dimensions. In: *Theoretical and Applied Aspects of Eye Movement Research*,
 765 edited by Gale AG, Johnson F. Amsterdam: Elsevier, 1984, p. 71–78.
- 766 Freedman, EG. Coordination of the eyes and head during visual orienting. *Exp Brain*
 767 *Res* 190(4): 369–387, 2008.

- 768 Fuller, JH. Head movement propensity. *Exp Brain Res* 92:152–164, 1992.
- 769 Gandhi NJ, Keller EL. Spatial distribution and discharge characteristics of superior
770 colliculus neurons antidromically activated from the omnipause region in the
771 monkey. *J Neurophysiol* 78: 2221–2225, 1997.
- 772 Groh, JM, Sparks, DL. Saccades to somatosensory targets: II. Motor convergence in
773 primate superior colliculus. *J Neurophysiol* 75: 428–438, 1996.
- 774 Goossens HH, Van Opstal AJ. Dynamic ensemble coding of saccades in the monkey
775 superior colliculus. *J Neurophysiol* 95:2326–41, 2006.
- 776 Guillaume A. Saccadic inhibition is accompanied by large and complex modulations
777 when induced by visual backward masking. *J Vis* 12: 1-20, 2012.
- 778 Hafed, ZM, Clark, JJ. Microsaccades as an overt measure of covert attention shifts.
779 *Vis Res* 42(22): 2533–2545, 2002.
- 780 Hafed, ZM, Ignashchenkova, A. On the Dissociation between Microsaccade Rate and
781 Direction after Peripheral Cues: Microsaccadic Inhibition Revisited. *J*
782 *Neurosci* 33(41): 16220–16235, 2013.
- 783 Hooge ITC, Erkelens CJ. Control of fixation during a simple search task. *Percept*
784 *Psychophys* 58: 969–976, 1996.
- 785 Jóhannesson, ÓI, Kristjánsson, Á. Violating the main sequence: asymmetries in
786 saccadic peak velocities for saccades into the temporal versus nasal
787 hemifields. *Exp Brain Res* 227(1): 101–110, 2013.
- 788 Keller, EL, Edelman, JA. Use of interrupted saccade paradigm to study spatial and
789 temporal dynamics of saccadic burst cells in superior colliculus in monkey. *J*
790 *Neurophysiol* 72(6): 2754–2770, 1994.

- 791 Ludwig CJH, Gilchrist ID, McSorley E. The remote distractor effect in saccade
792 programming: channel interactions and lateral inhibition. *Vis Res* 45: 1177–
793 1190, 2005.
- 794 Ludwig CJ, Mildinhall JW, Gilchrist ID. A population coding account for systematic
795 variation in saccadic dead time. *J Neurophysiol* 97: 795–805, 2007.
- 796 McIntosh, RD, Buonocore, A. Saccadic inhibition can cause the remote distractor
797 effect, but the remote distractor effect may not be a useful concept. *J Vis*,
798 14(5), 15–15, 2014.
- 799 Munoz DP, Wurtz RH. Saccade-related activity in monkey superior colliculus. I.
800 Characteristics of burst and buildup cells. *J Neurophysiol* 73: 2313–2333,
801 1995a.
- 802 Munoz DP, Waitzman DM, Wurtz RH. Activity of neurons in superior colliculus
803 during interrupted saccades. *J Neurophysiol* 75: 2562-2580, 1996.
- 804 Nuthmann A, Smith TJ, Engbert R, Henderson JM. CRISP: A computational model
805 of fixation durations in scene viewing. *Psychol Rev* 117: 382–405, 2010.
- 806 Olivier E, Dorris MC, Munoz DP. Lateral interactions in the superior colliculus, not
807 an extended fixation zone, can account for the remote distractor effects
808 (Commentary on Findlay & Walker). *Behavioral and Brain Sciences* 22: 694–
809 695, 1999.
- 810 Ottes FP, Van Gisbergen JA, Eggermont JJ. Visuomotor fields of the superior
811 colliculus: a quantitative model. *Vision Res* 26: 857–873, 1986.
- 812 Paré, M, Hanes, DP. Controlled movement processing: superior colliculus activity
813 associated with countermanded saccades. *J Neurosci* 23(16): 6480–6489,
814 2003.

- 815 Reichle E, Pollatsek A, Fisher D, Rayner K. Toward a model of eye movement
816 control in reading. *Psychol Rev* 105: 125–157, 1998.
- 817 Reingold EM, Stampe DM. Saccadic inhibition in complex visual tasks. In: *Current*
818 *Oculomotor Research: Physiological and Psychological Aspects*, edited by
819 Becker W, Deubel H, Mergner T. New York: Plenum Press, 1999 p. 249–255.
- 820 Reingold EM, Stampe DM. Saccadic inhibition and gaze contingent research
821 paradigms. In: *Reading as a Perceptual Process*, edited by Kennedy A,
822 Radach R, Heller D, Pynte J. Amsterdam: Elsevier, 2000, p. 119–145.
- 823 Reingold EM, Stampe DM. Saccadic inhibition in voluntary and reflexive saccades. *J*
824 *Cogn Neurosci* 14: 371–388, 2002.
- 825 Reingold EM, Stampe DM. Using the saccadic inhibition paradigm to investigate
826 saccadic control in reading. In: *The Mind's Eye: Cognitive and Applied*
827 *Aspects of Eye Movement Research*, edited by Hyona J, Radach R, Deubel H.
828 Amsterdam: Elsevier, 2003, p. 347–360.
- 829 Reingold EM, Stampe DM. Saccadic inhibition in reading. *J Exp Psychol Hum*
830 *Percept Perform* 30: 194–211, 2004.
- 831 Rizzolatti GG, Buchtel, HAH, Camarda, RR, Scandolara, CC. Neurons with complex
832 visual properties in the superior colliculus of the macaque monkey. *Exp Brain*
833 *Res* 38: 37–42, 1980.
- 834 Rolfs, M, Kliegl, R, Engbert, R. Toward a model of microsaccade generation: The
835 case of microsaccadic inhibition. *J Vis*, 8(11): 5–5, 2008.
- 836 Sparks, DL. The brainstem control of saccadic eye movements. *Nat Rev Neurosci*
837 3(12): 952–964, 2002.

- 838 Trappenberg, TP, Dorris, MC, Munoz, DP, Klein, RM. A Model of Saccade Initiation
839 Based on the Competitive Integration of Exogenous and Endogenous Signals
840 in the Superior Colliculus. *J Cognitive Neurosci* 13(2): 256–271, 2001.
- 841 Van Gisbergen JA, Van Opstal AJ, Tax AA. Collicular ensemble coding of saccades
842 based on vector summation. *Neuroscience* 21: 541–555, 1987.
- 843 Van Loon EM, Hooge ITC, Van den Berg AV. The timing of sequences of saccades
844 in visual search. *Proc R Soc Lond B Biol Sci* 269: 1571–1579, 2002.
- 845 Van Opstal, AJ, Goossens, HH. Linear ensemble-coding in midbrain superior
846 colliculus specifies the saccade kinematics. *Biol Cybern* 98: 561–577, 2008.
- 847 Walker R, Kentridge RW, Findlay JM. Independent contributions of the orienting of
848 attention, Fixation offset and bilateral stimulation on human saccadic latency.
849 *Exp Brain Res* 103: 294– 310, 1995.
- 850 Walker R, Deubel H, Schneider WX, Findlay JM. Effect of remote distractors on
851 saccade programming: evidence for an extended fixation zone. *J Neurophysiol*
852 78: 1108–1119, 1997.
- 853 Walshe RC, Nuthmann A. Mechanisms of saccadic decision making while encoding
854 naturalistic scenes. *J Vis* 15(5), 21–19, 2015.
- 855 Weber H, Aiple F, Fischer B, Latanov A. Dead zone for express saccades. *Exp Brain*
856 *Res* 89(1): 214–222, 1992.

857 **Figure captions.**

858

859 **Figure 1.** Experimental designs. Panel A. Trial sequence of Experiment 1.
 860 Participants were required to maintain fixation and to make a saccade to a red dot
 861 (0.5° of visual angle) appearing on the left or on the right side of the fixation cross at
 862 10° of eccentricity. Participants were also instructed to report if one out of the four
 863 briefly presented probes had a different orientation from the others (50% of trials) or
 864 if instead all probes were the same. On Flash trials, a black square (3.5° of visual
 865 angle) was presented at fixation for 11.7 ms in order to elicit SI. Participants were
 866 asked to ignore the flash. Panel B. Trial sequence of Experiment 2. Participants were
 867 instructed to maintain fixation and then to saccade toward a white stimulus presented
 868 on the left/right side of the screen at either 4, 8 or 12 degrees of visual angle. In Flash
 869 trials, two white bars were covering the 1/3 of the top and 1/3 of the bottom of the
 870 screen. Participants were asked to ignore the flash. Panel C. Trial sequence of
 871 Experiment 3. Similar structure of Experiment 2 but restricted to one target
 872 eccentricity (10 degrees). In Flash trials, a white bar could be presented either 4
 873 degrees less eccentric than target location (as showed in figure) or 4 degrees more
 874 eccentric than target location. In all figures stimuli are not in scale.

875

876 **Figure 2.** Example of saccadic main sequence fit. Three observers with similar main
 877 sequence values were chosen, one for each Experiment, and plotted with their
 878 respective fits. Panel A depicts Experiment 1, panel B Experiment 2 and pane C
 879 Experiment 3. The empty blue dot symbols show each observation in the no-flash
 880 trials while the full black dots represent velocities predicted from the corresponding
 881 amplitudes, based on the individual fit. Experiment 1 (panel A) and Experiment 3

(panel C) had only one eccentricity (10 degrees) and the fitted function is a 1st order polynomial. Experiment 2 (panel B) had a range of eccentricities (4, 8 and 12 degrees) and the fitted function is a 2nd order polynomial. R^2 for each experiment and participant are reported below.

Experiment 1: 0.15; 0.02; 0.61; 0.22; 0.09; 0.16; 0.05; 0.50; 0.20.

Experiment 2: 0.85; 0.75; 0.66; 0.70; 0.76; 0.76; 0.72; 0.84; 0.81; 0.92.

Experiment 3: 0.21; 0.02; 0.04; 0.51; 0.09; 0.25; 0.19; 0.02.

Figure 3. SI, saccadic gain and normalised peak speed in Experiment 1, 2 and 3. Panel A, D and G. Average SI profiles with 95% CI (shaded area). In Experiment 2, the SI profile was averaged also across the three eccentricities since we did not find any statistical difference among the three conditions. In Experiment 3, red represent flash- and green flash+ conditions (same convention in plot H and I). Variation in saccadic gain (panel B, E and H) and normalised peak speed (panel C, F and I) for target only (blue) and target plus flash (red and red/green in Experiment 3) trials. Data are binned in 20 ms intervals. Time on the x-axis is relative to flash onset; x-axis values thus represent the temporal lead of the flash relative to the observed launching of the saccade. Asterisks indicate significant differences between the no-flash and flash conditions (FDR corrected). Shaded areas represent the standard error of the mean.


902 **Authors contribution**

903

904 Antimo Buonocore and David Melcher designed Experiment 1, and all three authors
905 designed Experiments 2 and 3. Antimo Buonocore carried out data collection and data
906 analysis. Interpretation of the data was done equally by the three authors. Antimo
907 Buonocore drafted the manuscript and Robert McIntosh and David Melcher provided
908 critical revisions. All authors approved the final version of the manuscript for
909 submission.


A

Experiment 1




B




Experiment 2



C

Experiment 3
















Table 1. Maximum of SI (Dip maximum) and latency of the maximum (Dip latency) for each participant and condition in Experiment 1. Mean and S.D. are reported in the bottom row.

Experiment	Participant	Condition	Dip maximum	Dip latency
1	1	Central flash	0.90	81
	2		0.68	77
	3		0.84	85
	4		0.82	77
	5		0.64	73
	6		0.62	81
	7		0.97	97
	8		0.95	101
	9		0.61	97
	Mean		0.78	85
S.D.		0.14	10.3	

Table 2. Maximum of SI (Dip maximum) and latency of the maximum (Dip latency) for each participant and condition in Experiment 2. Mean and S.D. are reported in the bottom row.

Experiment	Participant	Condition	Dip maximum	Dip latency
2	1	Target 4°	0.85	81
	2		0.90	65
	3		0.70	77
	4		0.63	89
	5		0.64	69
	6		0.88	89
	7		0.73	73
	8		0.73	65
	9		0.62	93
	10		0.71	77
	Mean		0.74	78
	S.D.		0.10	10.1
2	1	Target 8°	0.82	89
	2		0.90	65
	3		0.60	109
	4		0.74	73
	5		0.50	81
	6		0.88	89
	7		0.79	73
	8		0.79	65
	9		0.60	77
	10		0.75	73
	Mean		0.74	79
	S.D.		0.13	13.4
2	1	Target 12°	0.96	85
	2		0.89	65
	3		0.53	109
	4		0.91	73
	5		0.68	73
	6		0.91	93
	7		0.85	69
	8		0.75	65
	9		0.55	65
	10		0.71	69
	Mean		0.77	77
	S.D.		0.15	14.7

Table 3. Maximum of SI (Dip maximum) and latency of the maximum (Dip latency) for each participant and condition in Experiment 3. Mean and S.D. are reported in the bottom row.

Experiment	Participant	Condition	Dip maximum	Dip latency
3	1	flash+	0.71	89
	2		0.29	93
	3		0.41	69
	4		0.66	93
	5		0.39	73
	6		0.36	73
	7		0.86	81
	8		0.57	73
	Mean		0.52	74
	S.D.		0.23	22.5
3	1	flash-	0.91	85
	2		0.81	73
	3		0.69	69
	4		1.00	89
	5		0.64	69
	6		0.78	73
	7		0.97	77
	8		0.88	81
	Mean		0.74	71
	S.D.		0.23	18.4